



UNIVERSITY of
ROCHESTER
MEDICAL CENTER

Neuromuscular Laboratory
601 Elmwood Avenue Room 5-5329
Rochester NY, 14621
585 275-1330

Skin Biopsy Instructions for referral skins processed at the University of Rochester.

Purpose: The purpose of this procedure is to ensure that every skin biopsy stained for Small Fiber Neuropathy testing at the University of Rochester is of maximum quality.

Scope: This procedure is for thick skin samples stained in the Neuromuscular Laboratory. It is performed on every referral skin biopsy and results are reported in the appropriate Anatomic Pathology Report.

Ordering Skin Biopsy Kits:

Skin biopsy kits can be ordered Monday through Friday 8 AM until 4:30 PM by calling the Neuromuscular Pathology Laboratory at 585-275-1330 or emailing at NMD_Pathology@urmc.rochester.edu. We will provide kits to all outside referring physicians that will include all the necessary tubes and solutions. **These kits have a shelf life of one year from preparation.** The tubes for the thigh will be labeled proximal thigh. The lateral distal leg tubes will be labeled distal leg/ankle. Tools for performing the biopsy will also be included.

Punch Skin Biopsy:

A standard biopsy to look for a distal small fiber sensory neuropathy includes a total of two 3-mm punch biopsies – one performed on the upper lateral thigh, and one performed on the lateral distal leg, about 7-10 cm above the lateral malleolus. **The tubes for the thigh will be labeled proximal thigh. The lateral distal leg tubes will be labeled distal leg/ankle.** Note that skin biopsies will be performed on the same side as the pain and/or burning. Additional skin biopsies can be taken from more proximal sites, or the contralateral side depending on the clinical presentation.

Important considerations in performing skin biopsy for nerve evaluation:

Is the person allergic to lidocaine?

Is the person on anticoagulants or have a coagulopathy?

Does the patient have a latex allergy?

Skin Biopsy Procedure:

The preparation consists of a thorough cleansing with 70% isopropyl alcohol.

Inject 1% lidocaine anesthetic with epinephrine (omit epinephrine if patients have had prior reactions to the epinephrine) is then infiltrated under the skin to raise a small bleb.

After the area is anesthetized, an Acupunch 3-mm punch is used to gently pull the skin plug up and away from the underlying subcutaneous fatty tissue. Punches from Accuderm only need to penetrate half the distance of the metal head. Deeper will only create excessive bleeding and reach fat layers.

Forceps should **NEVER** contact the upper layers of the skin. Handle gently by the deeper reticular dermis **ONLY**. These fine unmyelinated epidermal fibers will be destroyed if crush artifact is introduced.

A scalpel is used to cut any tissue ties.

Each specimen is then placed immediately in labeled specimen containers using Zamboni's fixative (tube A) for the anti PGP9.5 immunostaining and quantitation of the epidermal nerve fibers.

The site should then be dressed with a gauze pressure dressing.

Notes:

Only use Zamboni fixative provided. Formalin blocks the antigenic determinant sites. Fixation time should be 12-24 hours for optimal immunohistochemistry results.

Specimens must never be allowed to air dry on gauze. They should be placed directly in fixative.

When placing specimens in refrigerator, check to see that they are submerged in fixative and not stuck to the underside of the lid.

In determining side of subject to biopsy for peripheral neuropathy evaluation, do not biopsy distal to past surgical sites (i.e. past knee surgery). Do not biopsy skin

1% lidocaine injection should be close to the surface in order to raise a bleb. If injected only deep below the skin, the subject will still feel pain.

Stay away from lidocaine needle track when selecting punch site. Stay within bleb area but not including hole from needle.

Place a 4x4 inch gauze pad over biopsy site to prevent potential blood leakage.

You may batch ship fixed cryoprotected samples on wet ice/ice packs after holding in the refrigerator.

You must contact the lab before sending specimens. Specimens should not be shipped to arrive on a holiday or on weekends.

If you have questions regarding these protocols, please call the Neuromuscular Pathology laboratory at (585) 275-1330.

Sample Fixation and Preparation for Shipping:

This is a very critical step as proper fixation is essential for proper staining and interpretation.

The specimens should be placed in Tube A (Zamboni fixative) as soon as they are removed from the patient.

The specimens should remain in Tube A for 12-24 hours. The samples should be stored in the refrigerator during this time. If the samples stay in fixative for longer than 24 hours the results of the test will be compromised.

The specimens should be transferred to Tube B (Saline Solution) for 15 minutes.

The specimens should then be placed in Tube C (Shipping Buffer).

Samples should be stored in the refrigerator until they can be shipped to Rochester.

Shipping Information:

Specimens must be sent overnight: **Monday through Wednesday** to:

Neuromuscular Pathology Laboratory
University of Rochester Medical Center
Dept. of Neurology Rm. 5-5329
575 Elmwood Avenue
Rochester NY 14642

Contact Information:

E-mail NMD_pathology@urmc.rochester.edu
FAX: (585) 273-1255
Phone: (585) 275-1330

